

Discussion of paper LESA: Longitudinal Elastic Shape Analysis of Brain Subcortical Structures

Moo K. Chung¹, Jamie L. Hanson², Richard J. Davidson¹, Seth D. Pollak¹ *
Department of Biostatistics and Medical Informatics
¹University of Wisconsin-Madison, ² University of Pittsburgh

The brain surfaces including both cortical and subcortical structures including hippocampus have been analyzed for more than a decade using publicly available software packages such as FreeSurfer (Dale & Fischl 1999) and SurfStat (Worsley et al. 2009, Chung et al. 2010). Zhang et al. (2022) proposes an elastic shape metric based method for performing longitudinal shape analysis on brain subcortical structures. However, the demonstrated applications are limited to global summary measures such as the total surface area and principle component (PC) scores significantly limiting the impact of the study. For analyzing total surface area, we do not even need to align structures using LESA. PC scores lose richer vertex-based local information and it is unclear what parts of the hippocampus are responsible for longitudinal change. A more effective approach is to perform local shape analysis using the deformation-based morphometry (DBM) and tensor-based morphometry (TBM) after obtaining deformation in LESA (Ashburner et al. 1998, 2000, Thompson et al. 2000). Considering elastic methods put severe constraints on the Jacobian determinant of image deformation (Chung et al. 2001), it is not clear LESA can be effectively used in local shape analysis. We contrast shape analysis done in Zhang et al. (2022) against DBM and TBM in a longitudinal hippocampus study (Chung et al. 2011).

The *deformation-based morphometry* (DBM) utilizes the deformation field obtained

*The authors gratefully acknowledge support of grants NIH R01 EB028753 and NSF MDS-2010778.




Figure 1: Hippocampus longitudinal study of 124 children showing localized growth pattern difference between high-income (above \$75000) and low-income (below \$35000) families (Chung et al. 2011). The F -statistics map on testing the interaction between income and age while controlling for sex in a mixed-effects model is computed in SurfStat. The arrows are the average displacement differences between high and low income families. The posterior region is enlarging while the midbody and the anterior parts of right hippocampus are shrinking in low-income families (corrected p -value < 0.03). The developmental pattern is the opposite for high-income families.

through nonlinear image registration (Ashburner et al. 1998, Chung et al. 2001). In DBM, it is possible to detect local shape differences within the hippocampus and identify exactly

what subregion of hippocampus is responsible for the most growth (Figure 1). Given surface \mathcal{M} , the *deformation* is given as a 3D vector field $d(x)$ at vertex $x \in \mathcal{M}$. The deformation can be represented in the Lagrangian coordinate system as $d(x) = x + U(x)$, where $U = (U_1, U_2, U_3)$ is the *displacement* in the elastic deformation theory and measures a relative movement of vertex x (Chung et al. 2001). The longitudinal change over time t can be then modeled as $\frac{\partial U}{\partial t}(x, t) = L(U) + \Sigma^{1/2}(x)\epsilon(x)$. If the change is assumed to follow a diffusive behavior, then L is the Laplacian. If the morphological changes follow a fluid dynamics model or elastic deformation, L becomes the Navier-Stokes or elastic operator (Chung et al. 2001). Σ is the symmetric positive definitive covariance matrix allowing correlations between components of the deformations. The error vector field ϵ is assumed to be zero mean and unit standard deviation possibly Gaussian (Worsley et al. 1996).

In contrast to DBM, TBM quantifies the differential qualities called the displacement tensor $\frac{\partial U}{\partial x} = (\frac{\partial U_i}{\partial x_j})$ (Thompson et al. 2000, Chung et al. 2001). Suppose surface \mathcal{M} is parameterized by $x = X(v)$ with parameters $v = (v^1, v^2) \in \mathbb{R}^2$. The partial derivatives $X_i = \partial X / \partial v^i$ forms the basis in the tangent space. The Riemannian metric tensor $g = (g_{ij})$ is then given by the inner product $g_{ij} = \langle X_i, X_j \rangle$, which measures the amount of the deviation from the flat Euclidean plane. Unlike analyzing the total surface area $\int_{X^{-1}(\mathcal{M})} \sqrt{\det g} du$ as in Zhang et al. (2022), we can analyze the local *area element* $\sqrt{\det g}$ at the vertex resolution, the generalization of the Jacobian determinant, often used in TBM (Chung et al. 2003). Often DBM and TBM provide complimentary local shape information (Chung et al. 2001).

Since Zhang et al. (2022) only analyzed single summary measure per surface, it does not have multiple comparisons. The multiple comparisons across all the vertices is traditionally handled through the random field theory (Worsley et al. 1996). The random field theory is implemented in most brain imaging tools such as SPM (Ashburner et al. 1998) and

SurfStat (<https://laplcebeltrami.github.io/SurfStat>) (Worsley et al. 2009, Chung et al. 2010). SurfStat is the most widely used MATLAB package for building both fixed- and mixed-effects models for brain surface data. Mixed-effect model parameters are estimated using the restricted maximum likelihood (REML). SurfStat utilizes a model formula approach similar to R and we can simply set up a mixed-effect model used in Figure 1 as

```
lm = 1+ Sex + Age + Group + Age*Group + random(Subject) + I;
```

making variable `Subject` into a random effect. Identity matrix `I` is added to allow for independent noise in every subject. Then the corrected corrected p -value of test statistic T is reported as $P(\sup_{x \in \mathcal{M}} T(\mathbf{x}) \geq y)$. Figure 1 displays the output of SurfStat.

References

- Ashburner, J., Good, C. & Friston, K. (2000), ‘Tensor based morphometry’, *NeuroImage* **11S**, 465.
- Ashburner, J., Hutton, C., Frackowiak, R. S. J., Johnsrude, I., Price, C. & Friston, K. J. (1998), ‘Identifying global anatomical differences: deformation-based morphometry’, *Human Brain Mapping* **6**, 348–357.
- Chung, M., Hanson, J., Davidson, R. & Pollak, S. (2011), ‘Effect of family income on hippocampus growth: Longitudinal study’, *17th Annual Meeting of the Organization for Human Brain Mapping* **2697**. <https://www.stat.wisc.edu/~mchung/papers/HBM2011/chung.2011.HBM.pdf>.
- Chung, M., Worsley, K., Brendon, M., Dalton, K. & Davidson, R. (2010), ‘General multivariate linear modeling of surface shapes using SurfStat’, *NeuroImage* **53**, 491–505.

- Chung, M., Worsley, K., Paus, T., Cherif, D., Collins, C., Giedd, J., Rapoport, J. & Evans, A. (2001), ‘A unified statistical approach to deformation-based morphometry’, *NeuroImage* **14**, 595–606.
- Chung, M., Worsley, K., Robbins, S., Paus, T., Taylor, J., Giedd, J., Rapoport, J. & Evans, A. (2003), ‘Deformation-based surface morphometry applied to gray matter deformation’, *NeuroImage* **18**, 198–213.
- Dale, A. & Fischl, B. (1999), ‘Cortical surface-based analysis I. Segmentation and surface reconstruction’, *NeuroImage* **9**, 179–194.
- Thompson, P., Giedd, J., Woods, R., MacDonald, D., Evans, A. & Toga, A. (2000), ‘Growth patterns in the developing human brain detected using continuum-mechanical tensor mapping’, *Nature* **404**, 190–193.
- Worsley, K., Marrett, S., Neelin, P., Vandal, A., Friston, K. & Evans, A. (1996), ‘A unified statistical approach for determining significant signals in images of cerebral activation’, *Human Brain Mapping* **4**, 58–73.
- Worsley, K., Taylor, J., Carbonell, F., Chung, M., Duerden, E., Bernhardt, B., Lyttelton, O., Boucher, M. & Evans, A. (2009), ‘SurfStat: A MATLAB toolbox for the statistical analysis of univariate and multivariate surface and volumetric data using linear mixed effects models and random field theory’, *NeuroImage* **47**, S102.
- Zhang, Z., Wu, Y., Xiong, D., Ibrahim, J., Srivastava, A. & Zhu, H. (2022), ‘LESA: Longitudinal elastic shape analysis of brain subcortical structures’, *Journal of the American Statistical Association* (just-accepted), 1–27.