

Discussion 6

Review – Inference for One Population Mean

Point estimator: a natural point estimator for population mean is the sample mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$, with standard error = s/\sqrt{n} , where

$$s = \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}}.$$

sampling distribution of \bar{X} : By Central Limit Theorem, \bar{X} is approximately normal when n is large, even if we do not know the underlying distribution of population.

The following results are based on the assumption that the underlying distribution of population is *normal*.

P% confidence interval for μ :

$$\bar{x} - t^* \frac{s}{\sqrt{n}} < \mu < \bar{x} + t^* \frac{s}{\sqrt{n}},$$

where t^* is the critical value such that $P(|T_{n-1}| < t^*) = P\%$. Here, T_{n-1} denotes a random variable having t-distribution with degrees of freedom = $n-1$.

In R, `qt(p,df)` gives the quantile q such that $P(T_{df} < q) = p$.

Hypothesis Test:

- choose proper H_0 and H_A .

$$H_0 : \mu = \mu_0 \quad H_A : \mu \neq \mu_0 \text{ or } \mu < \mu_0 \text{ or } \mu > \mu_0.$$

- test statistic

$$T = \frac{\bar{x} - \mu}{s/\sqrt{n}}.$$

In R, the function `t.test()` carries out the t-test.

Hypothesis Test:

- choose proper H_0 and H_A .

$$H_0 : \mu = \mu_0 \quad H_A : \mu \neq \mu_0 \text{ or } \mu < \mu_0 \text{ or } \mu > \mu_0.$$

- test statistic

$$T = \frac{\bar{x} - \mu}{s/\sqrt{n}}.$$

In R, the function `t.test()` carries out the t-test.

Practice Problem

- Let $Z \sim N(0, 1)$. Find the following:
 - $P(Z > 2)$
 - $P(-1.96 < Z < 1.96)$
 - the constant c such that $P(|Z| < c) = 0.9$.
 - if $Y \sim N(10, 6)$, what is the distribution of $\frac{Y-10}{6}$?
- The mean height of women in Britain is 163.3 cm with standard deviation 6.4 cm. Let \bar{X} be the mean height of 36 British women randomly chosen from the population. Find:
 - Find $P(160 < \bar{X} < 168)$.
 - Find 0.95 quantile of the sampling distribution of \bar{X} .
 - Find the cutoff values of the middle 90% of the sampling distribution of \bar{X} .
- Six healthy three-year-old female Suffolk sheep were injected with the antibiotic Gentamicin, at a dosage of 10 mg/kg body weight. Their blood serum concentration of Gentamicin 1.5 hours after injection were as follows:

33 26 34 31 23 25

- Construct a 90% confidence interval for the population mean.
 - Interpret your result of (a)
 - The interval in (a) nearly contains all of the observations; will this typically be true for a 90% confidence interval? Explain.
 - Suppose we know that the s.d for the sheep's blood serum concentrations of Gentamicin 1.5 hours after injection is 5. How many samples do we need in order to guarantee that the error of estimation does not exceed 2?
- Madison Department of Transportation needs to analyze the average speed of cars passing downtown area, in order to see whether more traffic lights are needed for safety. Suppose, it is believed that an average speed of 15mph (or 24.14 kmph:kilometers per hour) is ideal for the balance of safety and smooth traffic. The velocity-detecting radar records 10 cars speed as shown in the following table. Use a two-sided test to analyze this dataset and represent your conclusions. (choose $\alpha = .05$)

| Cars | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | \bar{X} | S |
|--------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----------|-------|
| Speed (mph) | 13.31 | 13.59 | 16.22 | 19.13 | 17.42 | 22.25 | 18.30 | 22.32 | 17.34 | 16.97 | 17.685 | 3.047 |
| Speed (kmph) | 21.42 | 21.87 | 26.10 | 30.78 | 28.03 | 35.80 | 29.44 | 35.91 | 27.90 | 27.30 | 28.455 | 4.9 |

Solution

1.

```
> 1-pnorm(2)
[1] 0.02275013
> pnorm(1.96)-pnorm(-1.96)
[1] 0.9500042
> qnorm(0.05)
[1] -1.644854
```
2. By central limit theorem, \bar{X} will be approximately normal with mean 163.3 and $\text{sd } \frac{6.4}{\sqrt{36}} = 1.067$.
 - a. $P(160 < \bar{X} < 168) = P\left(\frac{160-163.3}{1.067} < \frac{\bar{X}-163.3}{1.067} < \frac{168-163.3}{1.067}\right) = P(-3.093 < Z < 4.405) = ?$

```
> pnorm(4.405)-pnorm(-3.093)
[1] 0.999004
```
 - b.

```
> qnorm(0.95,163.3,1.067)
[1] 165.0551
```
 - c.

```
> qnorm(c(0.05,0.95),163.3,1.067)
[1] 161.5449 165.0551
```
3. (a) $\bar{x} = 28.667$, $s = 4.59$, $n = 6$, $t^* = 2.015$. Then, the 90% confidence interval is (24.991, 32.442).
 - (b) Interpretation: We are 90% sure that the true mean blood serum concentration of Gentamicin is between 24.991 and 32.442.
 - (c) In general, this is not true for a 90% confidence interval. The meaning of a 90% confidence interval is not to say that such intervals will cover 90% data points in a sample.
 - (d) If the margin of error equals 2, we have

$$t_{n-1}^* \frac{s}{\sqrt{n}} \leq 2.$$

Hence,

$$\sqrt{n} \geq \frac{t_{n-1}^* s}{2}.$$

Note that, in the above formula, the quantile t_{n-1}^* also depends on n . Since there is no closed form for the quantile from t-distribution. We use the following R program to figure out the lower bound for n .

```
> s=5
> for (n in 2:50){
+   q=qt(0.95,n-1)
+   if(sqrt(n) > q*s/2) break;
+ }
> n
[1] 19
```

Therefore, $n = 19$ is already enough to reach a margin of error less than 2.

4.

$$H_0 : \mu = 15, \quad H_A : \mu \neq 15.$$
$$T = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} = \frac{17.685 - 15}{3.047/\sqrt{10}} = 2.787.$$

Then, the p-value equals

$$2 \times P(T_9 > 2.287) = 0.021 < \alpha.$$

Hence, we reject H_0 .