

Predicting Air Quality STAT 451 Final Project

Lacey Dinh, Tan Bui, Minji Suh, Maddie Young, and Diego Ugaz Why do we care about "fresh air"?

Motivation

- Air pollution can lead to illness such as lung disease, asthma, etc.
- Poor air quality results in 100,000 premature deaths and costs \$150 billion each year (National Weather Service)


Goal: investigate how meteorological and environmental factors influence air quality



Data Exploration

- Collected by N. Elgiriyewithana (<u>Kaggle</u>)
- Contains 40+ and over 37,000+ observations
- Key categories for features:
 - Time and Location: latitude, longitude, time of observation
 - Weather: temperature, precipitation, UV index
 - Pollutant: concentration of Nitrogen Oxide, Ozone
 - Astronomy: sunrise, moon phase

Research Questions



Preprocessing

- A subset of weather and pollutant features are used
 - Temperature, wind speed, humidity, cloud, UV index
 - Concentration of air pollutants
- No missing data!
- Variables are standardized to minimize the impact of outliers



Which factors play the biggest roles in determining the AQI level?

4 regression models:

- Linear Regression
- Decision Tree
- Random Forest
- Gradient Boosting

Evaluated based

on MSE and R^2

Results

Scores on Validation Set

DATA EXPLORATION

MSE

Optimal

 R^2

Hyperparameter

200 decision trees with a

max depth of 20

200 decision trees with an

alpha of 0.2

QUESTIONS

Model Linear Regression **Decision Tree**

Random Forest

Gradient Boosting

0.429 0.405



0.498





0.750

0.648




0.284

Q Results

Decision Tree:

Nodes near the root use concentrations of CO and SO₂ to branch.




Results

Random Forest:

Most important features for AQI are CO, SO_2 , NO_2

Scores on Test set:

- MSE: 0.201
- R^2 : 0.759




Q Results

Gradient Boosting:

Most important features for AQI are CO, SO₂, NO₂

Scores on Test set:

- MSE: 0.244
- R²: 0.708




- 0.2

- -0.2

Q1 Possible Complications

- Noticed similar importance of Ozone and Temperature in both ensemble models
- Ozone seems to be the most impactful pollutant on meteorological data
- Given these correlations, the meteorological features could be interpreted as repeat inclusions of Ozone as a feature






2 Do regions with similar climate patterns have similar air quality profiles?

Method:

K-means clustering

Features:

- Location & Weather variables
- Pollutants



2 Do regions with similar climate patterns have similar air quality profiles?

13


Cluster 2: Arid regions.

Cluster 4: Temperate

regions.

Cluster 6: Tropical

regions.




How do we classify AQI levels based on weather and pollutant data?

Method:

- Random Forest
- k-NN
- SVM
- Logistic Regression

Features:

- Weather variables
- Pollutants



How do we classify AQI levels based on weather and pollutant data?

Confusion Matrix (Test Set) - Random Forest 2172 Good -0 122 Moderate -15 0 Unhealthy for Sensitive Groups -19 0 555 6 0 Actual Unhealthy -24 70 24 1 Very Unhealthy -17 0 88 23 68 0 Hazardous -0 9 1 6 Unhealthy Good Unhealthy for Sensitive Groups Very Unhealthy Hazardous Moderate

Predicted

15

2000

1750

1500

1250

1000

750

500

- 250

- 0

Confusion Matrix:

AQI Documentation:

https://document.airnow.gov/tech nical-assistance-document-for-th e-reporting-of-daily-air-quailty .pdf

Conclusions and Limitations

16

Q1:

Most important features: carbon monoxide, sulfur dioxide, nitrogen dioxide.

Q2:

- Climate shapes air quality profiles across regions.
- Lack of temporal data limits insights into seasonal variations.

Q3:

- Skewed distribution toward "Good" AQI, reducing performance for rare categories.
- Model can aid in early warnings for air quality, improving public health response.

Thank You!

Let us know if you have any suggestions or questions