

# Localized Cortical Surface Asymmetry Analysis

Moo K. Chung, Kim Dalton, Richard J. Davidon  
Waisman Laboratory for Brain Imaging and Behavior  
University of Wisconsin-Madison

<http://www.stat.wisc.edu/~mchung>

## Abstract

*We present a computational framework for analyzing brain hemispheric asymmetry without any kind of image flipping. In order to perform brain asymmetry analysis, it is necessary to flip 3D magnetic resonance images (MRI) and establish the hemispheric correspondence by registering the original image to the flipped image. The difference between the original and the flipped images is then used as a measure of cerebral asymmetry. Instead of physically flipping MRI and performing image registration, we construct the global algebraic representation of cortical surface using spherical harmonics. Then using the inherent angular symmetry present in the spherical harmonics, image flipping is done by changing the sign of the asymmetric part in the representation. The surface registration between hemispheres and different subjects is done algebraically within the representation itself without any time consuming numerical optimization. The methodology has been applied in localizing the abnormal cortical asymmetry pattern of in a group of autistic subjects using the logistic discriminant analysis. Since the logistic discriminant analysis avoids the traditional hypothesis testing paradigm, the complicated multiple comparison issue that plagues the brain imaging community has been avoided.*

## 1. Introduction

Previous neuroanatomical studies on MRI have mainly flipped the whole brain 3D MRI to obtain the mirror reflected MRI with respect to the midsagittal cross section [2, 16]. The anatomical correspondence across the hemispheres is established and the normalized asymmetry index of type  $(L-R)/(L+R)$  is used at each voxel for quantification. Then the asymmetry index is feed into a statistical procedure, mainly general linear model (GLM), and a hypothesis on the estimated GLM parameters is tested and the resulting p-value is projected onto a template at each voxel. In order to address the correlated test statistic across voxels, multiple comparison procedures such as the random field

theory [29], false discovery rate (FDR) [4] and permutation tests [22] are utilized to construct the *corrected p-value* that account for multiple comparisons.

We present a radically different framework from this well established brain asymmetry analysis paradigm. Our proposed asymmetry analysis framework starts with segmenting the cortical surfaces using a deformable algorithm [blind] and obtaining cortical thickness that measures the distance between outer and inner cortical surfaces. Cortical thickness varies locally by region and is likely to be influenced by aging, development and clinical statue [3]. An algebraic representation of cortical surface is constructed using spherical harmonics. Inter-hemispheric and between-subject surface registrations are done within the algebraic representation without any numerical optimization that is needed for many previous surface registration techniques [8, 9, 21, 28]. The previous methods solve a complicated optimization problem of minimizing the measure of discrepancy between two surfaces while maximizing the smoothness of deformation. Our proposed technique does not try to normalize the original cortical meshes, which are highly noise, but instead normalize the algebraic representation of the cortical surfaces. Utilizing the property of Hilbert space, on which the algebraic representation resides, optimization is performed algebraically without the usual numerical optimization.

Instead of physically mirror reflecting the original MRI, the algebraic surface representation is mirror reflected by simply changing the sign in the representation. Then the inter-hemispheric correspondence can be established and the normalized asymmetry index on cortical thickness can be computed. The asymmetry index is feed into the logistic discriminant framework [13]. Unlike previous approaches [14, 24, 27] that classify preselected feature vectors, which are obtained mainly from the principal component analysis, our formulation can locally discriminate shape features at each mesh vertex. The methodology has been applied in localizing the abnormal cortical asymmetry pattern of in a group of autistic subjects and obtain the localized discriminant power up to 85.7%. The proposed method is simi-

lar to the traditional hypothesis driven statistical parametric mapping technique [29]. However, unlike the traditional approach, since there is no hypothesis to test, there is no p-value to report and there is no need to account for correlated test statistics across vertices. Hence the proposed discriminant analysis framework can completely bypass the complicated multiple comparison issue.

## 2. Methods

The human cerebral cortex has the topology of a 2D highly convoluted grey matter shell of average thickness of 3mm. The outer boundary of the shell is called the *outer cortical surface* while the inner boundary is called the *inner cortical surface*. Cortical surfaces are segmented from magnetic resonance images (MRI) using a deformable surface algorithm [blind] and mapped to unit sphere  $S^2$ . The resulting surfaces are represented as high resolution triangle meshes with the average inter-vertex distance of 3mm.

Let  $\zeta$  be the bijective mapping from  $\mathbf{u} = (u_j) \in S^2$  to the point  $\mathbf{p} = (p_j) \in \mathcal{M}$ , the outer cortical surface. The mapping  $\zeta$  enforces the one-to-one correspondence between the outer cortical mesh and the spherical mesh. The coordinates  $\mathbf{u}$  is parametrized by Euler angles  $(\theta, \varphi)$ :

$$(u_1, u_2, u_3) = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$$

with  $\Omega = (\theta, \varphi) \in \mathcal{N} = [0, \pi] \otimes [0, 2\pi)$ . The polar angle  $\theta$  is the angle from the north pole and the azimuthal angle  $\varphi$  is the angle along the horizontal cross section. We further parameterize surface  $\mathcal{M}$  with Euler angles in such a way that the plane  $u_2 = 0$ , which passes through  $\varphi = 0, \pi$ , is mapped to the midsagittal cross section of the brain (Figure 1):  $\mathbf{p}(\Omega) = \zeta \circ \mathbf{u}(\Omega)$ .

The coordinates  $\mathbf{p}(\Omega)$  are expected to be noisy. To filter out mesh noisy, we perform *diffusion-based surface smoothing* [1, 6, 19, 25, 26] [blind] on mesh vertices. Let  $\Delta$  be the spherical Laplacian defined as

$$\Delta = \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2}.$$

Consider the isotropic diffusion of the coordinates  $\mathbf{p}$ :

$$\frac{\partial f}{\partial \sigma} = \Delta f, \quad f(\Omega, \sigma = 0) = \mathbf{p}(\Omega). \quad (1)$$

The solution of (1) is given in terms of the integral transform called *heat kernel smoothing* [23] [blind] as

$$K_\sigma * \mathbf{p}(\Omega) = \int_{S^2} K_\sigma(\Omega, \Omega') \mathbf{p}(\Omega') d\mu(\Omega'), \quad (2)$$

where  $d\mu(\Omega) = \sin \theta d\theta d\varphi$  and  $K_\sigma$  is the heat kernel with bandwidth  $\sigma$ . Heat kernel smoothing (1) has the spectral representation. Consider eigenvalues  $\lambda$  and eigenfunctions




Figure 1. Cortical surface parameterization using Euler angles. The plane  $u_2 = 0$ , which passes through  $\varphi = 0, \pi$ , is mapped to the midsagittal cross section of the brain. The north ( $\theta = 0$ ) and the south ( $\theta = \pi$ ) poles are chosen along the midsagittal cross section.

$g$  satisfying  $\Delta g + \lambda g = 0$ . The eigenfunction  $Y_{lm}$  corresponding to the eigenvalue  $l(l+1)$  is called the *spherical harmonic* of degree  $l$  and order  $m$ . The explicit form of  $Y_{lm}$  is given in [blind] [7]. With respect to the inner product

$$\langle f, g \rangle = \int_{S^2} f(\Omega)g(\Omega) d\mu(\Omega),$$

$Y_{lm}$  form the orthonormal basis in  $L^2(S^2)$ , the space of square integrable functions. The  $L^2$ -norm is defined as  $\|f\| = \langle f, f \rangle^{1/2}$ . For a vector functional  $\mathbf{f} = (f_j)$ , the norm is similarly defined as  $\|\mathbf{f}\| = (\sum_j \|f_j\|^2)^{1/2}$ .

Using spherical harmonics, the heat kernel is written as

$$K_\sigma(\Omega, \Omega') = \sum_{l=0}^{\infty} \sum_{m=-l}^l e^{-l(l+1)\sigma} Y_{lm}(\Omega) Y_{lm}(\Omega'). \quad (3)$$

By substituting (3) into equation (2) and exchanging the integral with the summation, heat kernel smoothing (2) has the following spectral representation:

$$K_\sigma * \mathbf{p}(\Omega) = \sum_{l=0}^{\infty} \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm}(\Omega), \quad (4)$$

where  $\mathbf{p}_{lm} = \langle \mathbf{p}, Y_{lm} \rangle$  are the spherical harmonic coefficients. The finite expansion of (4) has the Hilbert space interpretation as stated in Theorem 1.

**Theorem 1** *Let the subspace*

$$\mathcal{H}_k = \left\{ \sum_{l=0}^k \sum_{m=-l}^l \beta_{lm} Y_{lm} : \beta_i \in \mathbb{R} \right\} \subset L^2(S^2),$$

*which is spanned by up to  $k$ -th degree spherical harmonics. Let  $h_0$  be the solution of (1). Then the closest function  $h$  in the subspace  $\mathcal{H}_k$  to  $h_0$  is given by*

$$\sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm}(\Omega) = \arg \min_{h \in \mathcal{H}_k} \|h - h_0\|.$$

Unlike previous literature [1, 6, 19, 25, 26] [blind] that solve surface diffusion numerically using the finite difference scheme, Theorem 1 provides a new framework for performing surface diffusion by computing a series expansion. The advantage of this new framework is the ability to explicitly model surface diffusion using the Karhunen-Loeve expansion of a random field [17]. The Fourier coefficient vector  $\mathbf{p}_{lm}$  can be modeled to follow independent multivariate normal  $N(\mu_{lm}, D)$ , where the covariance matrix is of the form  $D = \text{diag}(\sigma_{l1}^2, \sigma_{l2}^2, \sigma_{l3}^2)$ . Within the same degree, equal covariance is assumed. This model assumption is equivalent to the following.

$$\begin{aligned} & \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm} \\ &= \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mu_{lm} Y_{lm} \\ &+ \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \epsilon_{lm} Y_{lm}, \end{aligned}$$

where  $\epsilon_{lm}$  are independent multivariate normal  $\sim N(0, D)$ . The components of the second term are the form of the Karhunen-Loeve expansion [17]. The normality assumption was tested using the Jarque-Bera statistic [15] for the parameters ( $k = 42, \sigma = 0.001$ ) used in the study. For instance, 10 out of total  $(42 + 1)^2 = 1,849$  Fourier coefficients of  $x$ -component in  $\mathbf{p}_{lm}$  do not show normality at significance level  $\alpha = 0.05$ . Since uncorrelated Gaussian random variables are independent, we have also computed the cross correlation of all 1,849<sup>2</sup> pairs of coefficients to check independence (Figure 2). The most pairs show very low correlation and the average correlation is 0.16 indicating the independence assumption is valid.

The finite series expansion given in Theorem 1 will be called as the *weighted Fourier series*. The weighted Fourier series shares similarity with the spherical harmonic representation [11, 12, 24] which has been used to model simpler anatomical objects such as hippocampus [24] and ventricles [11] in brain imaging. When  $\sigma = 0$ , the weighted Fourier series becomes exactly the spherical harmonic representation.

## 2.1. Model parameter estimation

There are many available methods for computing Fourier coefficients [blind] [5, 11, 12, 24]; however, these methods are not suitable for computing the Fourier coefficients for a cortical mesh due to fairly large number of vertices. The fast Fourier transform [5, 12] usually need a predefined regular grid so if the mesh topology is different for each cortical mesh, a time consuming interpolation is needed. The least squares method [11, 24] is also not suitable when the




Figure 2. Cross correlation of up to 42 degrees. Each rows and columns are arranged in a vectorized fashion starting with  $(l, m) = (0, 0), (1, -1), (1, 0), (1, 1), \dots$ . Right figure is the enlargement of small white square in the top left corner of the left figure. The average correlation is 0.16 and the most of correlations are extremely low indicating the independence of Fourier coefficients.

number of mesh vertices and the degree of the expansion are large.

Consider a functional data  $f$  in  $L^2(S^2)$ . The functional data  $f$  is measured at the finite number of points  $\Omega_1, \dots, \Omega_n$ . Then the weighted Fourier series representation evaluated at  $\Omega_j$  is

$$f(\Omega_j) = \sum_{l=0}^k \sum_{m=-l}^l e^{-\lambda(l+1)\sigma} f_{lm} Y_{lm}(\Omega_j). \quad (5)$$

The system of equations (5) can be written in a matrix form. Let

$$\begin{aligned} \mathbf{f} &= (f(\Omega_1), \dots, f(\Omega_n))', \\ \beta &= (f_{00}, f_{1,-1}, f_{1,0}, f_{1,1}, \dots, f_{kk})', \\ \mathbf{Y} &= (e^{-l(l+1)\sigma} Y_{lm}(\Omega_j)). \end{aligned}$$

Matrix  $\mathbf{Y}$  is the size of  $n \times (k + 1)^2$ . The columns of  $\mathbf{Y}$  are  $Y_{00}, Y_{1,-1}, Y_{1,0}, Y_{1,1}, \dots$  evaluated at  $\Omega_1, \dots, \Omega_n$ . Then the system of equations (5) can be written as  $\mathbf{f} = \mathbf{Y}\beta$ , where the Fourier coefficients are estimated as

$$\hat{\beta} = (\mathbf{Y}'\mathbf{Y})^{-1} \mathbf{Y}'\mathbf{f}. \quad (6)$$

For most cortical surface segmentation algorithms [9, 18],  $n > 40,000$  and the matrix size can easily reach the physical memory limit of the most personal computers for  $k \geq 42$ . To address this issue, we have developed an iterative method for solving a large least squares problem by decomposing it into smaller least squares problems [blind]. The algorithm shares similarity to the *matching pursuit method* [20] in the underlying Hilbert space construction but differs in numerical implementation and motivation. While our algorithm was developed to avoid the computational burden of inverting a huge linear equation, the matching pursuit method was developed to compactly decompose a time frequency signal into a linear combination of pre-selected pool of basis functions.




Figure 3. Cortical thickness is simulated from the sample cortical thickness. The ground truth is analytically constructed from the simulation. Then the weighted Fourier series representation are compared against the ground truth. The mean relative error is at 0.0012.

We briefly describe the algorithm. Initially, we estimate the first coefficient  $f_{00}$  by solving

$$f(\Omega_j) = f_{00}Y_{00}(\Omega_j).$$

If we let  $\widehat{f_{00}}$  to be the least squares estimation, the next set of coefficients  $f_{1,-1}, f_{1,0}, f_{1,1}$  are estimated by solving

$$f(\Omega_j) - \widehat{f_{00}}Y_{00}(\Omega_j) = \sum_{m=-1}^1 e^{-1(1+1)\sigma} f_{1m}Y_{1m}(\Omega_j).$$

The subsequent set of coefficients  $f_{2,-2}, \dots, f_{2,2}$  are estimated by solving

$$f(\Omega_j) - \sum_{l=0}^1 \sum_{m=-l}^l \widehat{f_{lm}}Y_{lm}(\Omega_j) = \sum_{m=-2}^2 e^{-2(2+1)\sigma} f_{2m}Y_{2m}(\Omega_j).$$

At each iteration, the residual of the previous fit is used to estimate the next set of coefficients. The process continues until the residual is no longer statistically significant. At the  $l$ -th iteration, the iterative algorithm inverts manageable smaller  $(2l-1) \times (2l-1)$  matrix instead of huge  $(k+1)^2 \times (k+1)^2$  matrix in (6).

## 2.2. Validation

The weighed Fourier series representation is validated against analytically constructed ground truth. We have used the cortical thickness of a subject in constructing the analytical ground truth. Consider a surface measurement of the form

$$g(\Omega) = \sum_{l=0}^k \sum_{m=-l}^l \beta_{lm}Y_{lm}(\Omega) \quad (7)$$

for given  $\beta_{lm}$ . Heat kernel smoothing of  $g$  is given as an exact analytic form, which serves as the ground truth for validation:

$$K_\sigma * g(\Omega) = \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \beta_{lm}Y_{lm}(\Omega). \quad (8)$$

Using the sample cortical thickness data, we simulated the measurement of the form (7) by computing  $\beta_{lm} = \langle g, Y_{lm} \rangle$  (Figure 3 top left). Then we have compared the weighted Fourier series representation of  $g$  against the analytical ground truth (8) along the surface mesh (Figure 3). We have used  $\sigma = 0.001$  and the corresponding optimal degree  $k = 42$ . The relative error is up to 0.013 at a certain vertex and the mean relative error over all mesh vertices is 0.0012. Our validation results demonstrates that the numerical implementation of the weighted Fourier series representation is sufficiently good.

## 2.3. Surface registration in function space

Comparing measurements defined at different cortical surfaces is not a trivial task due to the fact that no two cortical surfaces are identically shaped. We need to establish surface correspondence between hemispheres and between subjects for subsequent statistical analysis. We present a new surface registration technique utilizing the Hilbert space property of weighted Fourier series. Consider parameterizations  $\widehat{\mathbf{p}}$  and  $\widehat{\mathbf{q}}$  obtained from coordinates  $\mathbf{p}$  and  $\mathbf{q}$  respectively:

$$\widehat{\mathbf{p}}(\Omega) = \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm}Y_{lm}(\Omega),$$

$$\widehat{\mathbf{q}}(\Omega) = \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{q}_{lm}Y_{lm}(\Omega).$$

Suppose the surface  $\widehat{\mathbf{p}}$  is deformed to  $\widehat{\mathbf{p}} + \mathbf{d}$ , where  $\mathbf{d} = (d_j)$  is the displacement vector field to be estimated. We find optimal displacement  $\mathbf{d}$  that minimizes the discrepancy between  $\widehat{\mathbf{p}} + \mathbf{d}$  and  $\widehat{\mathbf{q}}$  in the finite subspace  $\mathcal{H}_k$  using the  $L^2$ -norm:

**Theorem 2** *Given parameterization  $\widehat{\mathbf{p}}$  and  $\widehat{\mathbf{q}}$ , the optimal displacement from  $\widehat{\mathbf{p}}$  to  $\widehat{\mathbf{q}}$  is*

$$\sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} (\mathbf{q}_{lm} - \mathbf{p}_{lm})Y_{lm} = \arg \min_{d_j \in \mathcal{H}_k} \|\widehat{\mathbf{p}} + \mathbf{d} - \widehat{\mathbf{q}}\|.$$

Theorem 2 shows that the optimal displacement is obtained by taking the difference between two parametric representations and matching coefficients of the same degree and order. Note that we are not taking the difference between the original noisy surface meshes. Theorem 2 can be further used to establish the inter-hemispheric correspondence by letting  $\widehat{\mathbf{q}}$  to be the mirror reflection of  $\widehat{\mathbf{p}}$ .

**Theorem 3** *If  $\widehat{\mathbf{p}}^*$  denotes the mirror reflection of  $\widehat{\mathbf{p}}$  with respect to the midsagittal cross section, the optimal displacement from  $\widehat{\mathbf{p}}$  to  $\widehat{\mathbf{p}}^*$  is*

$$-2 \sum_{l=0}^k \sum_{m=0}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm}Y_{lm} = \arg \min_{d_j \in \mathcal{H}_k} \|\widehat{\mathbf{p}} + \mathbf{d} - \widehat{\mathbf{p}}^*\|.$$




Figure 4. Demonstration of surface registration based on Theorem 2 with parameters  $k = 78$  and  $\sigma = 0.0001$ . The first surface  $\hat{\mathbf{p}}$  is registered to the last surface  $\hat{\mathbf{q}}$ . Each surface corresponds to  $\hat{\mathbf{p}} + \alpha \mathbf{d}$  for  $\alpha = 0, 0.2, 0.4, 0.6, 0.8, 1$ . The last surface is  $\hat{\mathbf{q}} = \hat{\mathbf{p}} + \mathbf{d}$ .

*Proof.* From the way the spherical coordinates are set up (Figure 1), we obtain

$$\begin{aligned} \hat{\mathbf{p}}^*(\theta, \varphi) &= \hat{\mathbf{p}}(\theta, 2\pi - \varphi) \\ &= \sum_{l=0}^k \sum_{m=-l}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm}(\theta, 2\pi - \varphi) \\ &= \sum_{l=0}^k \sum_{m=-l}^{-1} e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm}(\theta, \varphi) \\ &\quad - \sum_{l=0}^k \sum_{m=0}^l e^{-l(l+1)\sigma} \mathbf{p}_{lm} Y_{lm}(\theta, \varphi). \end{aligned}$$

The last decomposition into negative and positive orders is obtained from the property of spherical harmonics:

$$Y_{lm}(\theta, 2\pi - \varphi) = \begin{cases} -Y_{lm}(\theta, \varphi), & -l \leq m \leq -1, \\ Y_{lm}(\theta, \varphi), & 0 \leq m \leq l. \end{cases}$$

Then letting  $\hat{\mathbf{q}} = \hat{\mathbf{p}}^*$ , we obtain the result.  $\square$

Theorem 3 shows that the optimal inter-hemispheric correspondence is obtained by matching the parameterization  $\hat{\mathbf{p}}(\theta, \varphi)$  to  $\hat{\mathbf{p}}(\theta, 2\pi - \varphi)$ . Again note that we are not trying to match the original mesh coordinates  $\mathbf{p}(\theta, \varphi)$  but their algebraic representations.

## 2.4. Localized asymmetry index

Theorem 3 enables us to establish the inter-hemispheric correspondence and, in turn, to construct cortical thickness based asymmetry index. Suppose  $\hat{g}$  is the weighted Fourier series representation of cortical thickness  $g$ . At each position  $\hat{\mathbf{p}}(\theta, \varphi)$ , we have cortical thickness  $\hat{g}(\theta, \varphi)$ . Then from Theorem 3, we match  $\hat{g}(\theta, \varphi)$  to  $\hat{g}(\theta, 2\pi - \varphi)$  for the hemispheric correspondence.

Comparing the expansion of  $\hat{g}(\theta, \varphi)$  and  $\hat{g}(\theta, 2\pi - \varphi)$ , we see that the negative order terms are invariant while the positive order terms change the sign. Hence, we decompose  $\hat{g}(\theta, \varphi)$  into symmetric  $S$  and antisymmetric  $A$  parts

as  $\hat{g}(\theta, \varphi) = S(\theta, \varphi) + A(\theta, \varphi)$ , where

$$\begin{aligned} S(\theta, \varphi) &= \frac{1}{2} [\hat{g}(\theta, \varphi) + \hat{g}(\theta, 2\pi - \varphi)] \\ &= \sum_{l=1}^k \sum_{m=-l}^{-1} e^{-l(l+1)\sigma} g_{lm} Y_{lm}(\theta, \varphi) \end{aligned}$$

and

$$\begin{aligned} A(\theta, \varphi) &= \frac{1}{2} [\hat{g}(\theta, \varphi) - \hat{g}(\theta, 2\pi - \varphi)] \\ &= \sum_{l=0}^k \sum_{m=0}^l e^{-l(l+1)\sigma} g_{lm} Y_{lm}(\theta, \varphi). \end{aligned}$$

Then the *normalized asymmetry index* of type (L-R)/(L+R) is given by

$$N(\theta, \varphi) = \frac{\sum_{l=1}^k \sum_{m=-l}^{-1} e^{-l(l+1)\sigma} g_{lm} Y_{lm}(\theta, \varphi)}{\sum_{l=0}^k \sum_{m=0}^l e^{-l(l+1)\sigma} g_{lm} Y_{lm}(\theta, \varphi)}.$$

The numerator is the sum of all negative orders while the denominator is the sum of all positive and the 0-th orders. The larger the value of the index, the larger the amount of asymmetry. Figure 5 shows the asymmetry index for 3 selective subjects.

## 2.5. Localized discriminant analysis

For each subject, its normalized asymmetry index  $N(\theta, \varphi)$  is computed and modeled as a zero mean Gaussian random field. For the traditional group comparison between autistic and normal control subjects,  $T$  statistic at each point  $(\theta, \varphi)$  would be constructed. Since  $T$  statistics at different points are correlated, it becomes a *multiple comparison problem* [4, 22, 29]. The corrected p-value of the test is determined by computing the superima distribution of the  $T$  random field [29], i.e.

$$P \left[ \sup_{(\theta, \varphi) \in S^2} T(\theta, \varphi) < h \right]. \quad (9)$$




Figure 5. Three representative subjects showing cortical thickness ( $g$ ), its weighted-SPHARM representation ( $\hat{g}$ ), asymmetry index ( $A$ ), symmetry index ( $S$ ) and normalized asymmetry index ( $N$ ). The Cortical thickness is projected onto the original brain surfaces while all other measurements are projected onto the 42-th degree weighed Fourier series representation.

Unfortunately, computing the suprime distribution of the  $T$  random field is not an easy task and requires satisfying many distributional assumptions and the estimation of the smoothness of the random field. We use a different approach called the *logistic discriminant analysis* [10, 13] that bypasses the multiple comparison issue and still able to detect the regions of abnormal asymmetry pattern in the autistic subjects. Unlike previous discriminant techniques [14, 24, 27] that tried to classify preselected feature vectors, our approach does not require any preselected feature vectors and performs the classification at each mesh vertex.

Let  $n_i$  denotes the normalized asymmetry index for the  $i$ -th subject at a particular point  $(\theta, \varphi)$ . Let  $Y_i$  be the clinical state of the  $i$ -th subject modeled as a Bernoulli random variable with parameter  $\pi_i$ .  $Y_i = 1$  if the  $i$ -th subject is autistic with probability  $\pi_i$  while  $Y_i = 0$  if the subject is normal with probability  $1 - \pi_i$ . Then we have the following logistic model, which links the probability of clinical statue  $\pi_i$  to the asymmetry index  $n_i$ :

$$\log \frac{\pi_i}{1 - \pi_i} = \beta_0 + \beta_1 n_i. \quad (10)$$

The unknown parameters  $\beta = (\beta_0, \beta_1)'$  are estimated by maximizing the loglikelihood function  $L(\beta)$ . The loglikelihood function is given by

$$\log L(\beta) = \text{const.} + \sum_{i=1}^n y_i (\beta_0 + \beta_1 n_i) + \log(1 - \pi_i). \quad (11)$$

Since the loglikelihood function (11) is not easy to maximize analytically, the Newton-Raphson method is used to maximize it in an iterative fashion. Starting with an arbitrary initial vector  $\beta^0$ , we estimate iteratively

$$\beta^{j+1} = \beta^j + I(\beta^j)^{-1} \frac{\partial \log L(\beta)}{\partial \beta}(\beta^j),$$

where  $I$  is the Fisher information matrix [10, 13].

Once we estimated the parameters, we classify the  $i$ -th subject as autistic if  $P(Y_i = 1) > P(Y_i = 0)$ , which is equivalent to the condition  $\pi_i > 1/2$ . The classification error rate  $\gamma(\theta, \varphi)$  at each point  $(\theta, \varphi)$  is estimated by the leave-one-out cross-validation scheme. Denote  $e_{-i}$  as the error rate for leaving the  $i$ -th subject out. Note that  $e_{-i} = 0$  if the subject is classified correctly while  $e_{-i} = 1$  if the subject is misclassified. Then the error rate  $\gamma$  is estimated as

$$\hat{\gamma} = \frac{1}{n} \sum_{i=1}^n e_{-i}.$$

The *discriminant power* is then given as  $1 - \hat{\gamma}$  and it is displayed in Figure 6 localizing the regions of abnormal asymmetry pattern in autistic subjects.

### 3. Application

Three Tesla  $T_1$ -weighted MR scans were acquired for 16 high functioning autistic and 12 control right handed males.

The autistic subjects were diagnosed by a trained and certified psychologist [blind]. The average ages are  $17.1 \pm 2.8$  and  $16.1 \pm 4.5$  for control and autistic groups respectively. The standard image processing steps, such as the intensity non-uniformity correction and the global affine normalization into the Montreal neurological institute stereotaxic space, were performed [blind]. Afterwards, an automatic tissue-segmentation algorithm based on a supervised artificial neural network classifier was used to segment gray and white matters. Cortical surface meshes were constructed by a deformable surface algorithm and cortical thickness  $g$  and mesh vertices  $\mathbf{p}$  are obtained. Using the bijective mapping from the unit sphere to the cortical surface, mesh vertices  $\mathbf{p}$  were parameterized by Euler angles  $(\theta, \varphi)$  (Figure 1).

The weighted Fourier series representation  $\hat{\mathbf{p}}$  and  $\hat{g}$  for 28 subjects was constructed using the iterative algorithm with bandwidth  $\sigma = 0.001$  corresponding to  $k = 42$  degrees (Figure 5). The representation has been validated against the ground truth and shown to perform sufficiently with the average relative error of 0.0012 (Figure 3). The symmetry ( $S$ ), asymmetry ( $A$ ) and normalized asymmetry ( $N$ ) indices were computed and projected onto  $k = 42$  degree representation (Figure 5). The normalized asymmetry index was used in localizing the regions of cortical asymmetry difference between two groups.

Instead of performing the usual two sample t-test, which introduces multiple comparison issues [4, 22, 29], logistic discriminant analysis was performed. At each point, the logistic model (10) was fitted to link the probability of clinical status to the asymmetry index. The logistic model was used to estimate the probability of the asymmetry index belonging to the autistic group. Then we computed the discriminant power, which is defined as the rate of correct classification. The discriminant power was projected onto the average cortical surface constructed by averaging the Fourier coefficients of all subjects within the same spherical harmonics basis using Theorem 2 (Figure 6). The average surface serves as an anatomical landmark for displaying these indices as well as projecting the logistic discriminant analysis result. The regions of high discriminant power indicates the likelihood of the regions to exhibit abnormal asymmetry in the autistic subjects.

## 4. Conclusions

We have presented a novel cortical asymmetry analysis technique called the weighted Fourier series representation that performs parameterization, smoothing and registration in a unified algebraic framework. The weighed Fourier series representation is given as the least squares approximation to an isotropic heat diffusion on a unit sphere. The diffusion time controls the amount of smoothing in the representation and enables the multiscale representation. The proposed method was able to connect diverse concepts of




Figure 6. Discriminant power projected on top of the average cortical surface. The discriminant power ranges from 32.1 to 85.7%. The logistic discriminant analysis framework provides an alternative to the traditional corrected p-value approach in the two group comparison setting and avoids complicated multiple comparison issue.

heat diffusion, surface registration and spherical harmonic representations in a consistent framework.

As an illustration, the methodology was applied quantifying the abnormal cortical asymmetry pattern of autistic subjects using the normalized asymmetry index. The asymmetry index that measures the amount of asymmetry presented in the cortical surface, was constructed as the ratio of spherical harmonics of negative and positive orders. The regions of abnormal pattern in the autistic subjects were localized using the logistic discriminant analysis. This completely avoids using the p-value formulation that accounts the multiple comparison issues using the random field theory [29].

## References

- [1] A. Andrade, F Kherif, J. Mangin, K.J. Worsley, A. Paradis, O. Simon, S. Dehaene, D. Le Bihan, and J-B. Poline. Detection of fmri activation using cortical surface mapping. *Human Brain Mapping*, 12:79–93, 2001. [2](#), [3](#)
- [2] T.R. Barrick, C.E. Mackay, S. Prima, F. maes, D. Vandermeulen, T.J. Crow, and N. Roberts. Automatic analysis of cerebral asymmetry: an exploratory study of the relationship between brain torque and planum temporale asymmetry. *NeuroImage*, 24:678–691, 2005. [1](#)
- [3] P. Barta, M.I. Miller, and A. Qiu. A stochastic model for studying the laminar structure of cortex from mri. *IEEE Transactions on Medical Imaging*, 24:728–742, 2005. [1](#)
- [4] Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B*, 57:289–300, 1995. [1](#), [6](#), [7](#)

- [5] T. Bulow. Spherical diffusion for 3d surface smoothing. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 26:1650–1654, 2004. 3
- [6] A. Cachia, J.-F. Mangin, Rivi re D., D. Papadopoulos-Orfanos, F. Kherif, I. Bloch, and J. R gis. A generic framework for parcellation of the cortical surface into gyri using geodesic vorono  diagrams. *Image Analysis*, 7:403–416, 2003. 2, 3
- [7] R. Courant and D. Hilbert. *Methods of Mathematical Physics: Volume II*. Interscience, New York, english edition, 1953. 2
- [8] C. Davatzikos. Spatial transformation and registration of brain images using elastically deformable models. *Comput. Vis. Image Underst.*, 66:207–222, 1997. 1
- [9] B. Fischl, M.I. Sereno, R. Tootell, and A.M. Dale. High-resolution intersubject averaging and a coordinate system for the cortical surface. *Hum. Brain Mapping*, 8:272–284, 1999. 1, 3
- [10] B. Flury. *A First Course in Multivariate Statistics*. Springer, 1997. 6
- [11] G. Gerig, M. Styner, D. Jones, D. Weinberger, and J. Lieberman. Shape analysis of brain ventricles using spharm. In *MMBIA*, pages 171–178, 2001. 3
- [12] X. Gu, Y.L. Wang, T.F. Chan, T.M. Thompson, and Yau. S.T. Genus zero surface conformal mapping and its application to brain surface mapping. *IEEE Transactions on Medical Imaging*, 23:1–10, 2004. 3
- [13] T. Hastie, R. Tibshirani, and J. Friedman. *The elements of statistical learning*. Springer, 2003. 1, 6
- [14] R. Higdon, N.L. Foster, R.A. Koeppe, C.S. DeCarli, W.J. Jagust, C.M. Clark, N.R. Barbas, S.E. Arnold, J.L. Heidebrink R.S. Turner, and S. Minoshima. A comparison of classification methods for differentiating fronto-temporal dementia from alzheimer’s disease using fdg-pet imaging. *Stat Med*, 23:315–326, 2004. 1, 6
- [15] C.M. Jarque and A.K. Bera. A test for normality of observations and regression residuals, journal = International Statistical Review, year = 1987, volume = 55, pages = 1-10,. 3
- [16] D.N. Kennedy, K.M. O’Craven, B.S. Ticho, A.M. Goldstein, N. Makris, and J.W. Henson. Structural and functional brain asymmetries in human situs inversus totalis. *Neurology*, 53:1260–1265, 1999. 1
- [17] J. Li and A.O. Hero. A spectral approach to statistical polar shape modeling. In *Proceedings of International Conference on Image Processing*, 2002. 3
- [18] J.D. MacDonald, N. Kabani, D. Avis, and A.C. Evans. Automated 3-d extraction of inner and outer surfaces of cerebral cortex from mri. *NeuroImage*, 12:340–356, 2000. 3
- [19] R. Malladi and I. Ravve. Fast difference schemes for edge enhancing beltrami flow. In *Proceedings of Computer Vision-ECCV*, pages Part I 343–357, 2002. 2, 3
- [20] S. Mallat and Z. Zhang. Matching pursuits with time-frequency dictionaries. *IEEE Transactions on Signal Processing*, 41:3397–3415, 1993. 4
- [21] M.I. Miller, A. Banerjee, G.E. Christensen, S.C. Joshi, N. Khaneja, U. Grenander, and L. Matejic. Statistical methods in computational anatomy. *Statistical Methods in Medical Research*, 6:267–299, 1997. 1
- [22] T. Nichols and S. Hayasaka. Controlling the family-wise error rate in functional neuroimaging: a comparative review. *Stat Methods Med. Res.*, 12:419–446, 2003. 1, 6, 7
- [23] S. Rosenberg. *The Laplacian on a Riemannian Manifold*. Cambridge University Press, 1997. 2
- [24] L. Shen, J. Ford, F. Makedon, and A. Saykin. surface-based approach for classification of 3d neuroanatomical structures. *Intelligent Data Analysis*, 8:519–542, 2004. 1, 3, 6
- [25] N. Sochen, R. Kimmel, and R. Malladi. A general framework for low level vision. *IEEE Transactions on Image Processing*, 7:310–318, 1998. 2, 3
- [26] B. Tang, G. Sapiro, and V. Caselles. Direction diffusion. In *The Proceedings of the Seventh IEEE International Conference on Computer Vision*. 2, 3
- [27] C. Thomaz, J.P. Boardman, S.J. Counsell, D.L.G. Hill, J.V. Hajnal, A.D. Edwards, M.A. Rutherford, D. Gillies, and D. Rueckert. A whole brain morphometric analysis of changes associated with preterm birth. In *SPIE Medical Imaging 2006: Image Processing*, volume 6144, pages 1903–1910, 2006. 1, 6
- [28] P.M. Thompson and A.W. Toga. A surface-based technique for warping 3-dimensional images of the brain. *IEEE Transactions on Medical Imaging*, 15, 1996. 1
- [29] K.J. Worsley, S. Marrett, P. Neelin, A.C. Vandal, K.J. Friston, and A.C. Evans. A unified statistical approach for determining significant signals in images of cerebral activation. *Human Brain Mapping*, 4:58–73, 1996. 1, 2, 6, 7